

Read This First: Quick Start Guide

+1 (888) 707-4355
sales@aviancontrol.com
aviancontrol.com

Avian Control®: What to Expect

Avian Control® utilizes food-grade Methyl Anthranilate (MA), which harmlessly irritates birds' taste and smell receptors upon contact (via inhalation, preening, ingestion, or even through their feet). This unpleasant sensation causes birds to leave the treated area. Unlike temporary fixes or scare tactics, birds never habituate to Avian Control®, ensuring reliable, ongoing bird reduction without harm to birds, staff, tenants, or property.

Training Period (1-4 Weeks)

- **Frequency:** Apply 1-2 times weekly during this initial phase, ensuring even coverage across the area birds are frequenting.
- **Birds May Still Visit:** It's normal to see birds after the first few applications. Like any training process, it isn't always an instant fix. Stick with it – the results are worth the patience.
- **Consistency is Crucial:** Regular, consistent application is key to effectively training the birds and reinforcing the message, especially for any newcomers exploring the area.



Once the birds learn to stay away, you can usually switch to a lower frequency – we recommend 2-4 monthly applications, but observe bird activity and adjust as needed.



Unlike scare tactics birds can get used to, they don't acclimate to Avian Control. Sticking with maintenance sprays is key to reinforcing the message, especially for any new birds exploring the area.

Application Tips

Rate For Efficacy

Do not use below recommended dosage, typically 32 oz. of product per acre, especially for initial treatments.

Apply as a Fine Mist

Ensure the target area receives thorough, even coverage. Run off will have no beneficial effect.

Reapply After Heavy Rain

Re-treat the area after each significant rainfall (1-2in.), as these can reduce effectiveness.



Read This First: Mixing Guide

Targeting Problem Areas

For Crops

Apply to the entire plant and fruit acreage, not just the perimeter. Start applications 1 month prior to growing season or at the first signs of bird presence.

Outdoor Fogging

Apply Avian Control® undiluted at a rate of 12 to 16 ounces per acre (43,500 sq ft). Use a dry fog with droplet size below 30 microns. Point fog generator in the general direction of birds, maintaining at least a 5-foot distance from structures/objects.

For Buildings/Structures

Apply directly to rooftops, ledges, HVAC units, and other structural surfaces where birds roost or nest.

Indoor Fogging

Apply Avian Control® undiluted at a rate not to exceed two (2) ounces per 10,000 cubic feet of space. Point the fog generator in the general direction of birds, keeping it at least 5 feet away from structures to allow fog dissipation. Thermal fogging is ideal for reaching high places as the hot fog naturally rises.

Mixing Chart

Important: Do not use Avian Control® with water that has a pH lower than 7. (common with well water). Add a buffer to increase pH to 7 or above.

	Application Area	Avian Control®	Water	InvisiDye™ (Optional)
Residential	¼ Acre (approx.)	8 oz	Fill to 1 gallon mark	½ oz
Commercial	½ Acre (approx.)	16 oz	Fill to 2 gallon mark	1 oz
Agriculture	1 Acre or Greater	32 oz per Acre	Enough water for full coverage	Not For Use