



Avian Enterprises, LLC
2000 Pontiac Drive
Sylvan Lake, Michigan 48320
Phone: 888-707-4355
Fax: 888-877-4355
sales@AvianControl.com
www.AvianControl.com

Avian Control® Bird Repellent Best Practices Guidance for Dairy Barns

Best Practices for Application of Avian Control® Bird Repellent in Dairy Barns:

The following methods are recommended to maximize the effectiveness of Avian Control® in dairy barns.

- Mist or fog the repellent in the barn twice a day the first week during the early morning & near dusk hours when the birds are present. Mist or fog the repellent in the dairy barn twice the second week, with follow-up applications every week thereafter until the birds are gone. This usually requires three to four weeks of applications. A fine mist or fog is most effective.

If you do not own a fly spray machine, misting or fogging machines, we recommend the SOLO 451 gas powered backpack mister blower. We have tested this machine and found it to be quite effective. It is available in stores and online.

- Mist, fog or spray a solution of Avian Control® Bird Repellent directly to the surface of the feedstuffs* two or three times daily in feed aisles, bunkers and other open areas. Spraying is recommended in the early morning hours to deter birds from eating the feed. This action should be done on a daily basis for the first two weeks along with misting or fogging the barn as described above. ***Note:** California and New York do not allow application of repellent to animal feedstuffs.
- Avian Control® Bird Repellent works best when diluted in neutral to slightly alkaline water. If using well water that is acidic, care should be given to neutralizing the water before use.

We welcome your comments and are always available for questions by phone or email. Please feel free to consult our website: www.aviancontrol.com for additional useful information.