## **READ THIS FIRST** Tips for Agricultural Users

- Always read and follow all label directions.
- Avian Control<sup>®</sup> Bird Repellent is typically applied at from 24 oz. to 32 oz. per acre using your standard volume of water. A surfactant may be used with Avian Control<sup>®</sup> if the spray system does not agitate the solution.
- Avian Control<sup>®</sup> Bird Repellent may be applied by air. Consult the label for specific instructions.
- Re-apply Avian Control® Bird Repellent at 7 to 10 day intervals.
- Avoid tank mixing Avian Control<sup>®</sup> Bird Repellent with low pH pesticides. Solutions with a pH of less than 7 (neutral) reduces its effectiveness. If your water has a pH lower than 7, add a buffer to increase the pH to 7.
- Avian Control<sup>®</sup> Bird Repellent may be used on listed crops up to the day of harvest. See the label for a complete list.
- Avian Control<sup>®</sup> Bird Repellent has an REI of 4 hours.
- Avian Control<sup>®</sup> Bird Repellent may be applied to non-avian feedstuffs.
- Avian Control<sup>®</sup> Bird Repellent may be used to remove birds from barns, packing sheds and other farm buildings.
- Avoid using Avian Control<sup>®</sup> Bird Repellent on dwellings such as homes, cabins, and other occupied buildings.



2000 Pontiac Dr., Sylvan Lake, MI 48320 • sales@AvianControl.com • www.AvianControl.com